Te Whatu Ora
Health New Zealand
Waikato

Patient name:

NHI:

Address:

Allergies:

Symptom Response Kit Medications (renal-impairment version)

Authority for subcutaneous administration

Infuse the following medications subcutaneously over 24 hours:

Drug	Dosage	Route
		CSCI
Water or 0.9% sodium chloride to volume.		

Increments:

	Increment increase / decrease amount	To a maximum dose of...	Frequency

Subcutaneous boluses:

Medication	Dosage	Frequency and indication
Fentanyl	25-50 mcg	Q1H prn for pain or dyspnoea
Midazolam	2.5 mg	Q1H prn for dyspnoea or anxiety
Haloperidol	0.5-1 mg	Q4H prn for nausea/vomiting or delirium/agitation
Buscopan (hyoscine butylbromide)	20 mg	Q2-4H prn for secretions, max 80mg/24h

Ensure prescriptions are provided for:

- Fentanyl 100mcg/2mL ampoules, supply 10 x 2mL ampoules
- Midazolam 15mg/3mL ampoules, supply 5 x 3mL ampoules
- Haloperidol 5mg/mL ampoules, supply 5 x 1mL ampoules
- Hyoscine butylbromide 20mg/mL ampoules, supply 10 x 1mL ampoule

Once completed, please email to clinical.admin@hospicewaikato.org.nz

Doctor's name and role:

MCNZ number: Date:

Doctor's signature: